

Galaxies and Superclusters

B.G. Sidharth*

Centre for Applicable Mathematics & Computer Sciences
B.M. Birla Science Centre, Adarsh Nagar, Hyderabad - 500 063 (India)

Abstract

In this brief note, we would like to point out that large-scale structures like galaxies and superclusters would arise quite naturally in the universe.

There being about $N = 10^{11}$ galaxies in the universe, each about $l = 10^{23} \text{cms}$ across, we can easily verify that

$$R \sim \sqrt{N}l \quad (1)$$

where R is the radius of the universe.

As is well known, (1) arises in the theory of Brownian motion on the one hand[1] and on the other, it is also true if N represents the number of elementary particles $\sim 10^{80}$, in the universe and l , their size or spread, that is their Compton wavelength[2].

We can now interpret (1) as follows:

From the large-scale perspective, the galaxies are approximately in Brownian motion and their size is given correctly by (1). Interestingly, as we can easily verify, an identical relation holds for superclusters also[3], with a similar interpretation.

Moreover (1) implies a two dimensional structure - this is indeed true; not only do galaxies have large flat disks, but also superclusters have a flat cellular character[4].

However, (1) is not true for stars. In this case gravitation is strong and the Brownian approximation is no longer valid.

Thus galaxies and superclusters would naturally arise in the Universe.

^{0*}Email:birlasc@hd1.vsnl.net.in; birlard@ap.nic.in

References

- [1] B.G. Sidharth, Int. J. Mod. Phys. A, 13 (15), 1998, p.2599ff.
- [2] F. Reif, "Fundamentals of Statistical and Thermal Physics", McGraw Hill, Singapore, 1965.
- [3] B.G. Sidharth, [xxx.lanl.gov/gr-qc 9903053](http://xxx.lanl.gov/gr-qc/9903053).
- [4] J.V. Narlikar, "Introduction to Cosmology", Cambridge University Press, Cambridge, 1993.